

Report: QPD 10.0
Patient ID: 12345678
Date: 1/1/2017 12:00:00 PM

Device: 1/1/2017 12:01:00 PM

Symptom Scores

Scale	Results		Reference Range
	within range	out of range	
Depression*		14	0 - 10
Manic Episode		5	0 - 3
Anxiety		14	0 - 10
Panic Disorder	6		0 - 8
PTSD		8	0 - 2
Eating Disorder	2		0 - 4
Substance Use	0		0 - 4
Somatization	5		0 - 11

* 11-14 mild / 15-19 moderate / 19-24 moderately severe / >24 severe

Diagnostic Notes:

- Patient appears to meet criteria for Persistent Depressive Disorder (Dysthymia)
- Possible Bipolar Mood Disorder, further inquiry indicated
- Clinically significant anxiety (does not meet formal criteria for Generalized Anxiety Disorder)
- Patient appears to meet criteria for PTSD

Report:	QPD 10.0 Report	Date Uploaded:	10/13/2015 12:01:00 PM
ID:	12345678	Name:	John Doe

Depression Symptoms

- depressed mood more days than not, 2 years or longer duration
- diminished interest or pleasure in activities
- weight loss
- insomnia
- fatigue, loss of energy
- feelings of worthlessness or guilt
- hopelessness

(Hypo)Manic Symptoms

The patient has answered 'true' to the following statements indicating a possible manic or hypomanic episode:

- There has been a period lasting four days or longer when I was excited, hyper, or "revved up" that people thought I was not myself, or I got myself into trouble.
- During this period, I needed much less sleep than usual.
- During this period, I talked too much without stopping, or talked so fast that people had trouble following me.
- During this period, my thoughts were racing through my head.
- During this period, I was so easily distracted by everything that it was hard to keep myself on one track.


Anxiety Symptoms

- excessive anxiety or worry
- worried about multiple things
- feels unable to stop worrying
- easily fatigued
- muscle tension
- sleep disturbance
- feels helpless and overwhelmed
- easily startled
- frequent heart racing or pounding
- frequent urination
- frequent dry mouth
- anxiety causes trembling or shaking

PTSD Symptoms

- intrusive memories of traumatic event
- distressing dreams of traumatic event
- flashbacks of traumatic event
- intense distress or physiological reactivity when reminded of traumatic event
- avoids stimuli associated with traumatic event
- emotional detachment or numbness
- feelings of being unable to live out a normal life
- insomnia
- startles easily

Depression and Anxiety Over Time



Disclaimer: Patient Tools, Inc. is a technology company that provides screening tools for development, mental health, behavioral health, and other issues. Patient Tools does not provide any medical, psychological, or professional services to its customers or users. For an accurate medical or mental health diagnosis, please see a qualified professional. Patient Tools employees, consultants and agents shall not be liable for any claims or damages, and expressly disclaim all liability of any nature for any action or non-action taken as a result of the information generated by the Patient Tools website or any of its programs. If you or a loved one is experiencing a medical emergency, please contact emergency services.